

# Application of Bacteriocins of *Lactobacillus brevis* as Biomedicine

Thapar Parul<sup>1\*</sup>, Malik R.K.<sup>2</sup> and Salooja M.K.<sup>3</sup>

1. CT University, Ludhiana, INDIA

2. National Dairy Research Institute, Karnal, INDIA

3. Indira Gandhi National Open University, New Delhi, INDIA

\*parul.thapar@yahoo.com

## Abstract

World Health Organization (W.H.O.) has pointed that despite great improvements in the oral health of population in several countries, dental caries still persists and has become a global problem. The oral health care demands are beyond the capacities of the health care systems in most low and middle income countries. Therefore, it was thought to explore the natural ways by which the problems of dental caries can be prevented in a cost-effective manner. The inhibition of caries causing organisms can be done by naturally occurring substances like bacteriocins found in lactic acid bacteria. In the present study, prevention of dental caries by utilization of bacteriocins from *Lactobacillus brevis* has been done by incorporating bacteriocin in toothpaste. Five strains of *Lactobacillus brevis* isolated from curd were used for the production of bacteriocin.

Out of five bacteriocin producers, the strain LB11(1) showed maximum percentage of MIC. The bacteriocin activity was optimized using different media, pH and temperature. The partial purification of the bacteriocin was done using ammonium sulphate precipitation and centrifugal filtration. The molecular weight of the purified bacteriocin was 3 kDa, while yield obtained was 100 per cent. The purified bacteriocin was then incorporated in the toothpaste to check the effect of inhibition on caries causing organisms (*Streptococcus mutans* and *Streptococcus sobrinus* isolated from dental cavities). It was confirmed that the bacteriocin induced toothpaste can be applied to prevent dental caries directly by the consumer and in a cost effective manner.

**Keywords:** Bacteriocins, lactic acid bacteria, *Streptococcus mutans*, *Streptococcus sobrinus*, *Lactobacillus brevis*, dental caries.

## Introduction

Dental caries is the most common disease occurring in all age groups which is preventable and is the major cause of oral pain and tooth loss<sup>13</sup>. Children suffering with tooth decay also suffer with improper growth and development which severely affect their health. India is a diverse country with different ethnic, geographic and cultural diversity.

Therefore, it is quite expected to have different disease patterns in various regions or states of the country<sup>11</sup>.

According to the report of World Health Organization, a large number of clinical treatments for dental caries have been developed with improved technologies in most of the countries of the world, but still the disease of dental caries has become a global burden. The oral health care demands are beyond the capacities of the health care systems in most low and middle income countries<sup>21</sup>. Keeping in view of the above data, it was thought to explore the natural ways by which the problems of dental caries can be prevented in a cost-effective manner.

It was found that a few studies have set the path through use of natural bacteriocins against inhibition of caries causing organisms. Bacteriocins are potent protein toxins produced by virtually every bacterial and archeal species<sup>7</sup> which can be applied to prevent dental cavities. Bacteriocins have a fast acting mechanism and they work even at extremely low concentrations. They form pores in the target membrane of bacteria, thus inhibiting the cell-wall biosynthesis<sup>16</sup>. Bacteriocins possess certain characteristics like narrow target range, stability, high activity and low toxicity. The bacteriocins of lactic acid bacteria can also be applied to food and medicine, where they can play a major role in preservation, packaging, food additives and prevention of certain diseases.

The objectives of the present study included: a) Isolation and screening of bacteriocins from the strains of *Lactobacillus brevis*. b) Partially purification of the bacteriocin and c) Incorporation of bacteriocin in toothpaste.

## Material and Methods

**Isolation of bacteriocin:** The five strains of *Lactobacillus brevis* [LB11(1), LB13(3), LB13(4), LB13(5), LB14(3)] isolated from curd were grown in De Man Rogosa and Sharpe (MRS) broth for 24 h at 24°C. Cells were removed by centrifugation at 12,000 rpm for 5 min in 1.5 ml sterilized Eppendorf tubes.<sup>5</sup> The culture supernatant thus obtained was transferred to another sterilized Eppendorf tube.

**Determination of anti-bacterial activity:** This supernatant (crude bacteriocin) was employed for determining the antibacterial activity. The indicator organisms used were ten strains of *Streptococcus mutans* [SM10(a), SM13(a), SM14(a), SM22(a), SM23(a), SM28(a), SM29(a), SM33(a), SM35(a), SM36(a)] and four strains of *Streptococcus sobrinus* [SS19(b), SS23(b), SS24(b), SS36(b)] (dental

caries causing organisms). These organisms were isolated from the dental cavities. The anti-bacterial activity was determined by Spot on Lawn method.

**Minimum Inhibitory Concentration (MIC) of the Crude Bacteriocin:** Minimum inhibitory concentrations (MICs) are defined as the lowest concentration of an antimicrobial that will inhibit the visible growth of a microorganism after overnight incubation<sup>3</sup>. To determine the MIC of the bacteriocins, the method as described in the Clinical and Laboratory Standards, 2010 was followed. Spot on lawn assay was done against the fresh indicator cultures grown at 24 h at optimum conditions.

Different concentrations (as given in the table 1) of the crude bacteriocin solution were put as drop on each plate of the isolate. The incubation was done at 37°C for 24 h. MIC was calculated as AU/ml= 200 x MIC. The statistical analysis of the MIC was performed using ANOVA- single factor.

**Growth Curve of Bacteriocin Producing Strain:** To determine the growth curve of bacteriocin producing strain, the method as described by the Hall et al<sup>9</sup> was followed. The selected bacteriocin producing cultures- LB11(1) was inoculated in sterilized De Man Rogosa and Sharpe (MRS) broth and incubated at 37°C. Optical density (O.D.) of the bacterial growth was measured at 600 nm after 2, 4, 6, 8, 10, 24, 30 and 36 h of incubation. To determine the maximum growth of the cells at a particular incubation hour, a growth curve was plotted against the time of incubation and O.D.

**Optimization of the Conditions for Maximum Production of Bacteriocin:**

- Medium- De Man Rogosa and Sharpe (MRS) and M-17 flasks inoculated with the cultures were incubated at 42°C.
- Initial pH- De Man Rogosa and Sharpe (MRS) broth adjusted to various initial pH values (5.0, 6.0 and 7.0) was inoculated with the cultures and incubated at 42°C.

- Temperature- De Man Rogosa and Sharpe (MRS) broth inoculated with the producer strain was incubated at 37°C, 42°C and 50°C.

In all the above experiments, the samples were drawn at 0, 8, 16, 24 and 36 h intervals and tested for pH and bacteriocin activity units. The results obtained were represented in the form of table and graphs. The statistical analysis for the parameter pH for all the conditions was statistically analyzed using ANOVA- single factor.

**Protein Estimation:** Protein content of the samples was estimated by the method of the Hartree,<sup>10</sup> Lowry et al<sup>12</sup> and Wilson and Walker<sup>20</sup> using bovine serum albumin (BSA) as the standard. The concentrations of the test solution were noted under absorbance at 600 nm which were analyzed by graphical representation. The amount of protein was calculated as:

$$\text{mg protein/ml} = \frac{\text{mg}(\text{protein}) (\text{Dilution})}{\text{Reagent} (\text{ml})}$$

**Ammonium Sulfate Precipitation:** The procedure as described by Yang et al<sup>22</sup> was followed. To 100 ml of supernatant, 37.32 g of ammonium sulfate was added slowly with constant stirring at 4- 5°C to achieve 60 per cent saturation and stirring continued for another 4 h at 5-7°C. The mixture was then kept overnight in refrigerator at 0°C. It was then centrifuged at 13,000 rpm for 20 min and the precipitates were dissolved in 0.1N phosphate buffer (pH 6.8). This partially purified bacteriocin was further subjected to purification by centrifugal filters.

**Centrifugal Filters:** The bacteriocin solution obtained was further purified using 3kDa MWCO (molecular weight cut off) centrifugal filter. The method as described by Gautam and Sharma<sup>8</sup> was used with some modifications. 0.5 ml of partially purified bacteriocin was added in two filters of 3kDa each in sterilized condition.

**Table 1**  
**Different concentrations**

S.N.	Concentration	CFCs	Distilled Water
1.	2x	100µl	100µl
2.	4x	100µl	300µl
3.	6x	100µl	500µl
4.	8x	100µl	700µl
5.	10x	100µl	900µl
6.	15x	100µl x (6)	50µl
7.	20x	100µl x (6)	100µl
8.	30x	100µl x (6)	200µl
9.	40x	100µl x (6)	300µl
10.	50x	50µl x (6)	450µl
11.	100x	100µl x (11)	100µl
12.	200x	100µl x (12)	100µl

These were centrifuged at 13,000 rpm for 10 min. The purified sample was recovered in another sterile Eppendorff tube. The specific activity and yield of the bacteriocin were calculated from the protein content and activity units of the samples. The Bacteriocin yield (%) was calculated as:

Weight of sample before filtration x 100/ Volume of sample used for filtration.

**Enzyme Sensitivity:** To know the sensitivity of the bacteriocin against enzyme, the proteolytic enzyme used was Protease. 5 ml of bacteriocin was treated with enzyme (1 mg/ml) at pH 7.0. Incubation was done at 37°C for 24 h. It was then heated at 100°C for 3- 5 min to denature the enzyme. This was then assayed by agar well-diffusion method for determining the antimicrobial activity. The standard method as described by the Barefoot and Klaenhammer<sup>4</sup> was followed.

The purified bacteriocin was incorporated in toothpaste (standard composition as mentioned in table 2 to determine the effect of these substituted products on caries causing organisms. Take 1g/1ml of toothpaste in four different sterile tubes and incorporate with different concentrations of bacteriocin- 0.05 ml, 0.1 ml, 0.5 ml and 1 ml.

Agar well assay was used to know the bacteriocin activity while being incorporated in the products. 50 µl of the incorporated bacteriocin were filled in agar wells on the solidified mat of indicator organism. These were incubated

at 37°C for 24 h to observe for the presence or absence of zone. A control sample of toothpaste without bacteriocin was also kept along for testing.

**Results and Discussion**


**A. Isolation and screening of bacteriocins from the strains of *Lactobacillus brevis***

**Anti-microbial assay against *Streptococcus mutans* and *Streptococcus sobrinus*:** The five strains of *Lactobacillus brevis* were used against the indicator organisms- nine strains of *Streptococcus mutans* and five strains of *Streptococcus sobrinus* to determine the antibacterial activity. All the five strains of *Lactobacillus brevis* showed zones of inhibition against six isolated strains of *Streptococcus mutans* and *Streptococcus sobrinus* species. as represented in the figure 1. The statistical analysis of the diameter of zone of inhibition was statistically analyzed using ANOVA. This showed that the maximum percentage of inhibition is shown by the isolate LB11(1)- 26% against the isolates of *Streptococcus mutans* and *Streptococcus sobrinus*.

All the values were average of three trials. The p-value within the isolates is 1.71E-09 and showed no significant difference at  $p \leq 0.05$ . In the figure 2, the graphical representation of the zone of inhibition has been presented showing the maximum inhibition by the isolate LB11(1) against the *Streptococcal* isolates.

**Table 2**  
**Standard composition of toothpaste**

S.N.	Constituents	Quantity
1.	Abrasives- aluminium hydroxide, calcium hydrogen phosphates, calcium carbonate, silica and hydroxyapatite	50.0 %
2.	Detergents- Sodium lauryl sulfate (SLS)	0.5-2.0 %
3.	Antibacterial agents- Triclosan or zinc chloride	30.0 %
4.	Sodium fluoride	1450 ppm
5.	Water	20- 40%
6.	Flavourants- spearmint, peppermint and wintergreen	1000 ppm



**a) Zone of inhibition formed against *Streptococcus mutans* and *Streptococcus sobrinus***  
**b) No zone of inhibition**

**Fig. 1: Anti-bacterial activity of the Strains against Streptococcal species**

The maximum inhibition shown by the strain LB11(1) may be due to the presence of optimal growth conditions for bacteriocin production.

The study by Ahmed et al<sup>2</sup> had also shown similar results as the diameter zones are included between 2 to 10 mm and the biggest clear zones were recorded by the extracts of *Pediococcus pentosaceus* NRC AM1 and *Pediococcus pentosaceus* NRC AM4 with *Escherichia coli* 0157:H7, but the lowest clear zones were obtained with the extracts of *Lactobacillus brevis* NRC AM2 and *Lactobacillus plantarum* NRC AM7 against *Pseudomonas aeruginosa*.

**Minimum Inhibitory Concentration (MIC) of the Crude Bacteriocin:** To determine the lowest concentration of an antimicrobial (bacteriocin), the Minimum Inhibitory Concentration (MIC) of the crude bacteriocin was determined by spot-on-lawn assay<sup>18</sup>. MIC was calculated as:

$$\text{AU/ml} = 200 \times \text{MIC}$$

The minimum inhibitory concentration was 0.02  $\mu\text{l}$  as shown by the strain LB11<sup>1</sup> with maximum antimicrobial activity of 10,000 AU/ml against the Streptococcal strain of SM28(a). The dilution of culture supernatant to 50X dilution did not show any zone of inhibition. The diameter of zone of inhibition obtained for all the strains was 0.1mm. The MIC of the crude bacteriocin was statistically analyzed using ANOVA-single factor. This represented the minimum inhibitory concentration as percentage and the maximum percentage of MIC is shown by bacteriocin producing strain LB11<sup>1</sup>- 2% against the isolates of *Streptococcus mutans* and *Streptococcus sobrinus*. All the values were average of three trials. The p-value within the strains is 0.002 and did not show any significant difference at  $p \leq 0.05$ .

The presence of optimal conditions required for the maximum activity of bacteriocin for the strain LB11<sup>1</sup> favored this strain to produce maximum percentage of inhibition at the minimum concentration.

In a study by the Agrawal and Prakash<sup>1</sup> similar results were shown. MIC of various isolates of lactic acid bacteria producing bacteriocins isolated from fermented milk was determined. The MIC ( $\mu\text{l}$ ) observed was as follows: *Lactococcus* spp.- 0.05, *Lactococcus* spp.- 0.6, *Leuconostoc*- 0.2, *Pediococcus* (4 different strains)- 0.6, 0.5, 0.5 and 0.05.

**Growth Curve:** To obtain the optimum growth conditions of the bacteriocin producing strain, a growth curve of isolated lactic acid bacterial strain LB11<sup>1</sup> in MRS broth was plotted against time of incubation and optical density (O.D.) at 600 nm which was incubated at 37°C for 36 h. It was observed that the maximum bacteriocin was produced at 24 h and production remained stable till 30 h. After reaching 36 h, the bacteriocin production started to decline and pH of MRS broth decreased from 5.97 to 3.6 (Figure 3). In a study by the Wang et al,<sup>19</sup> the values obtained were somewhat

similar- the growth curve of isolated strain of *Lactobacillus brevis* from Jiaoke (natural fermented cream in China) was plotted which showed that the maximum bacteriocin production was reached after 24 h and production remained stable from 24 h to the end of the growth. During 48 h of growth, pH of MRS broth decreased from 5.8 to approximately 3.6.

### Optimization of Conditions for Maximum Production of Bacteriocin:

- **Influence of Culture Medium on Bacteriocin Production:** The optimization of conditions for different media was done using De Man Rogosa and Sharpe (MRS) and M-17 media. A gradual decrease in the initial pH of the growth media (MRS and M17) was observed with increase in incubation period varying from 0-36 h. The bacteriocin production was observed maximum in MRS media at 24 h at pH 5.9 against the Streptococcal strain SM28(a) (Figure 4a).
- **Statistical analysis:** The optimization of conditions for different media for pH was statistically analyzed using ANOVA- single factor. The p-value within the strains for MRS media is 5.87E-15 and did not show any significant difference at  $p \leq 0.05$ . Similarly, the p-value within the strains for M-17 media is 2.56E-12 and did not show any significant difference at  $p \leq 0.05$ .

In a study by Namasivayam et al,<sup>15</sup> the activity of the bacteriocin produced from *Lactobacillus brevis* isolated from curd was evaluated against three bacterial species in MRS media supplemented with various nitrogen and carbon sources. MRS media supplemented with soybean meal recorded maximum anti- bacterial activity against the tested bacterial strains which is in support to the present study.

- **Influence of Initial pH on Bacteriocin Production:** The optimization of conditions for initial pH was done using pH 5.0, 6.0 and 7.0. De Man Rogosa and Sharpe (MRS) broth was adjusted to various initial pH values (5.0, 6.0 and 7.0) and inoculated with the cultures and incubated at 42°C. A gradual decrease in the initial pH range of 5.0, 6.0 and 7.0 was observed in the growth media at incubation period varying from 0-36 h. The bacteriocin activity was observed at 24 h and it showed decrease in the activity at 36 h in the pH of 5.0, 6.0 and 7.0. The bacteriocin production was shown to be maximum at pH 5.8 at 24 h against the Streptococcal strain SM28(a) (Figure 4b).
- **Statistical analysis:** The statistical analysis of the strains for comparison of the parameter pH was done using ANOVA- single factor. The p-value within the strains at pH 5.0 is 1.64E-06 and did not show any significant difference at  $p \leq 0.05$ . At pH 6.0, the p-value within the strains is 4.78E-13 and did not show any significant difference at  $p \leq 0.05$ . For

pH 7.0, the p-value within the strains is 1.84E-10 and did not show any significant difference at  $p \leq 0.05$ .

In a study, the bacteriocin strains of *Lactobacillus* species isolated from dairy products showed that all the strains were grown well at pH 4, 6 and 8 but only strains of *Pediococcus pentosaceus* NRCAM1, *Lactobacillus brevis* NRCAM2 and *Pediococcus pentosaceus* NRCAM4 were also grown at pH 2.0 which is in support to the present study.<sup>2</sup>

- Influence of Incubation Temperature on Bacteriocin Production:** The optimization of conditions has been done using different temperatures- 37°C, 42°C and 50°C. De Man Rogosa and Sharpe (MRS) broth inoculated with the culture was incubated at 37°C, 42°C

and 50°C. It was observed that with gradual rise in incubation period from 0 to 36 h, there was a gradual decrease in the initial pH. The rise in the incubation period from 0 to 36 h at 37°C showed maximum bacteriocin activity at 24 h which gradually decreased at 36 h of incubation whereas at 42°C and 50°C, there has been consistent decrease in the activity of bacteriocin. The maximum activity of bacteriocin was shown at 37°C and pH 5.9 for 24 h against the Streptococcal strain SM28(a).

Statistical analysis: Statistical analysis of the isolates for the parameter pH at different temperatures was performed. For temperature 37°C, the p-value within the strains is 3E-15 and did not show any significant difference at  $p \leq 0.05$ .




Figure 2: Percentage inhibition by strains




Fig. 3: Growth Curve




Fig. 4a: Different media




Fig. 4b: Different pH




Fig. 4c: Different temperature

At temperature 42°C, the p-value within the strains is 2.4E-14 and did not show any significant difference at  $p \leq 0.05$ . Also, for temperature 50°C, the p-value within the strains is 5.41E-15 and did not show any significant difference at  $p \leq 0.05$ .

In a study, the bacteriocin strains of *Lactobacillus* species isolated from dairy products including the strains of *Lactobacillus brevis* NRCAM2 showed the growth at 30°C and 37°C. On the other hand, strains of *Lactobacillus brevis* NRCAM2, *Lactococcus lactis* sub sp. *lactis* NRCAM3 and *Lactobacillus pentosaceus* NRCAM5 were not able to grow at temperature 45°C. On the other hand, the species of *Lactobacillus plantarum* and *Lactobacillus rhamnosus* can grow at 45°C which is in support to the present study<sup>2</sup>.

### B) Partial purification of the bacteriocin

The partial purification of the bacteriocin was done by using ammonium sulphate precipitation. The partially purified bacteriocin was passed through centrifugal filter of 3 kDa. After purification, the specific activity of the purified bacteriocin was determined by spot-on-lawn assay against the isolated strains of *Streptococcus mutans* and *Streptococcus sobrinus*. The activity units of the bacteriocin LB11(1) was 20,000 AU/ml against the isolated strain of *Streptococcus sobrinus*- SS19(b). The diameter of the zone of inhibition was 0.2 mm. The molecular weight cut-off of the purified bacteriocin after centrifugal filtration was 3kDa and the bacteriocin yield obtained was 100%.

A study by the Wang et al<sup>19</sup> showed similar results: the molecular weight of the bacteriocin of *Lactobacillus brevis* isolated from Jiaoke (natural fermented cream in China) was approximately 3.8 kDa according to tricine SDS-PAGE which is in support of the present study.

**Protein Estimation of the Crude and Partially Purified Bacteriocin:** The total protein content of the crude and purified bacteriocin of the strain LB11(1) was determined by

the method described by the Lowry et al<sup>12</sup>. The total protein content in crude bacteriocin was calculated as 1.0 mg/ml and that of purified bacteriocin was 0.2 mg/ml.


In a study by the Mohankumar and Murugalatha,<sup>14</sup> similar results were obtained which showed that the purified bacteriocin produced from the lactic acid bacteria had protein content of 0.21 mg/ml.

**Enzyme Sensitivity of Bacteriocin:** To determine the proteinaceous nature of the bacteriocin, the bacteriocin of the strain LB11(1) was treated with the enzyme- protease. It was observed that the bacteriocin had lost its inhibitory activity upon treatment with the proteolytic enzyme. The enzyme protease had cleaved the bacteriocin and this bacteriocin could not show its antimicrobial action against the indicator organism. This confirmed the proteinaceous nature of the bacteriocin.

The bacteriocin like substance produced by *Lactobacillus brevis* B23 showed no antimicrobial activity or it was unstable after heat treatment with all the proteolytic enzymes (protease and trypsin). The completely inactivated inhibitory activity indicated that it was proteinaceous in nature which is in support of the present study<sup>17</sup>.

### C) Incorporation of bacteriocin in toothpaste

Agar well assay was performed on solidified mat of indicator organism. The diluted concentrations for 100X and 200X of 50 µl toothpaste were poured into the wells of the medium and incubated at 37°C for 24 h. After incubation, these were observed for the presence or absence of zones. The zone of diameter at concentration of 100X for the control sample of toothpaste was 0.6 mm and the toothpaste with bacteriocin was 1.01 mm (average of three trials), while at concentration of 200X, the control sample of toothpaste was 0.8 mm and for the toothpaste with bacteriocin was 1.21 mm (average of three trials) (Figure 5).



\*\*WB: Without Bacteriocin

**Fig. 5: Increase in Inhibition of Bacteriocin Induced Toothpaste against Indicator Organism**






Fig. 6: Zone of diameter for inhibition by toothpaste

**Statistical analysis:** Statistical analysis of the zone of diameter for the bacteriocin induced toothpaste at concentration 100X and 200X was performed using ANOVA-single factor. The p-value within the isolates is  $8.16E-05$  and did not show any significant difference at  $p \leq 0.05$ . The comparison of zone of diameter for bacteriocin induced inhibition by toothpaste and control sample is shown in the figure 6.

## Conclusion

The five strains of *Lactobacillus brevis* isolated from curd were used for the production of bacteriocins. The crude bacteriocins obtained were applied to test the antimicrobial activity against the isolated Streptococcal species of dental cavities. All the five strains showed antimicrobial activity against the six indicator organisms of Streptococcal species. This indicated the presence of natural inhibiting substance called bacteriocin within the isolated *Lactobacillus brevis* strains. It was found that the minimum inhibitory concentration was obtained by the bacteriocin of the strain LB11(1)-0.02  $\mu$ l with maximum antimicrobial activity of 10,000 AU/ml against the Streptococcal strain of SM28(a) and LB11(1) was regarded as the best bacteriocin producing strain as the optimal condition for its growth was suitable. The statistical analysis revealed De Man Rogosa and Sharpe (MRS) to be superior to M-17 media at 24 h at pH 5.9 against the Streptococcal strain SM28(a).

This is because the MRS media contains all essential nutrients required for the bacteriocin production whereas there is absence of such nutrients in M-17 medium. The bacteriocin can be considered heat labile as the activity was decreased at the temperatures  $42^{\circ}\text{C}$  and  $50^{\circ}\text{C}$ . In all the conditions, there was a gradual decrease in the initial pH with gradual rise in incubation period from 0 to 36 h. This is due to the loss of activity due to nutrient limiting factor after 24 h or the bacteriocin is degraded by the enzymes. It was concluded that the bacteriocin producing strain LB11<sup>1</sup> showed maximum antimicrobial activity at  $37^{\circ}\text{C}$  for 24 h between pH 6.0-7.0.

The molecular weight cut-off of the purified bacteriocin after centrifugal filtration was 3kDa and the bacteriocin yield obtained was 100%. The total protein content in crude bacteriocin- 1.0 mg/ml and that of purified bacteriocin- 0.2 mg/ml. The proteinaceous nature of the bacteriocin was also confirmed by the application of enzyme protease. Thus, bacteriocins from *Lactobacillus brevis* can help in enhancing the prevention of dental caries as a biomedicine. This can be achieved by inoculating the bacteriocin in toothpaste as it can be applied directly by the consumers and patients of dental caries in an easy and cost-effective manner.

## References

1. Agrawal N. and Prakash A., Isolation of lactic acid bacteria from fermented milk products and their antimicrobial activity against *Staphylococcus aureus*, *J Food Safety*, **15**, 39-42 (2013)
2. Ahmed M.M., Baher A.M., Zainab I.M., Tawfik N.F. and Hassan Z.M.R., Antibacterial activity of some lactic acid bacteria isolated from Egyptian dairy products, *Int J Chem Tech Res*, **6**, 1139-1150 (2014)
3. Andrews J.M., Determination of minimum inhibitory concentrations, *J Antimicrob Chemother*, **48**, 5-16 (2002)
4. Barefoot S.F. and Klaenhammer T.R., Purification and characterization of *Lactobacillus acidophilus* bacteriocin lactacin B, *Antimicrob Agents Chemother*, **26**, 328-334 (1984)
5. Burlanek L.L. and Yousef A.E., Solvent extraction of bacteriocins from liquid cultures, *J Applied Microbiol*, **31**, 193-197 (2000)
6. Clinical and Laboratory Standards Institute: Performance standards for antimicrobial susceptibility testing; twentieth informational supplement, M100-S20 (2010)
7. Dorit L.R., Roy M.S. and Riley A.M., The bacteriocins current knowledge and future prospects book, Caister Academic Press (2016)
8. Gautam N. and Sharma N., Purification and characterization of bacteriocin produced by strain of *Lactobacillus brevis* MTCC 7539, *Ind J Biochem Biophy*, **46**, 337-341 (2009)



9. Hall B.G., Acar H., Nandipati A. and Barlow M., Growth rates made easy, *J Molecular Biol Evol*, **31**, 232-238 (2013)
10. Hartree E.E., Determination of protein: a modification of the lowry method that gives a linear photometric response, *J Anal Biochem*, **48**, 422 (1972)
11. Janakiram C., Antony B., Joseph J. and Ramanarayanan V., Prevalence of dental caries in India among the WHO index age groups: a meta- analysis, *J Clin Diag Res*, **12**, 8-13 (2018)
12. Lowry O.H., Rosebrough N.J., Farr A.L. and Randall R.J., Protein measurement with the Folin phenol reagent, *J Biol Chem*, **193**, 265 (1951)
13. Mathews S. and Mathew G.V., War inside the mouth, *J Dent Med Sci*, **13**, 23-25 (2014)
14. Mohankumar M. and Murugalatha M., Characterization and antibacterial activity of bacteriocin producing *Lactobacillus* isolated from raw cattle milk sample, *Int J Biol*, **3**, 128-143 (2011)
15. Namasivayam S.K.R., Angel J.C.R., Bharani R.S.A. and Karthik M.Y., Effect of Media on bacteriocin production by *Lactobacillus brevis* and evaluation of anti-bacterial activity, *Res J Pharm Biol Chem Sci*, **5**, 1129-1136 (2014)
16. Rodney H.P., Takeshi Z. and Kenji S., Novel bacteriocins from lactic acid bacteria (LAB): various structures and applications, *Microbial Cell Fact*, **13**, 1-3 (2014)
17. Rushdy A.A. and Gomaa E.Z., Antimicrobial compounds produced by probiotic *Lactobacillus brevis* isolated from dairy products, *Ann Microbiol*, **63**, 81-90 (2013)
18. Toro C.R., Use of probiotic lactic bacteria in feeding *Litopenaeus vannamei* shrimp as inhibitors of pathogenic microorganisms and immune system stimulants, Ph.D. Thesis, Federal University of Paraná, Curitiba, Brazil (2005)
19. Wang H., Gong X., Meng L. and Man L., A Bacteriocin Produced by *Lactobacillus brevis* KLDS1.0373 isolated from “Jiaoke”, a traditional fermented cream from China, *Advanced Materials Res*, **14**, 183-185 (2011).
20. Wilson K. and Walker J., Practical Biochemistry: Principles and Techniques, Cambridge University Press (2000)
21. World Health Organization, A report on Oral Health, from <https://www.who.int/news-room/fact-sheets/detail/oral-health> (2018)
22. Yang R., Johnson M.C. and Ray B., Novel method to extract large amounts of bacteriocins from lactic acid bacteria, *J Appl Environ Microbiol*, **58**, 3355-3359 (1992).

(Received 18<sup>th</sup> May 2020, accepted 25<sup>th</sup> July 2020)